

A mass generation mechanism for gauge fields

A. Sevostyanov

Department of Mathematical Sciences

University of Aberdeen

Aberdeen AB24 3UE, United Kingdom

e-mail: seva@maths.abdn.ac.uk

phone: +44-1224-272755, fax: +44-1224-272607

March 10, 2008

Abstract

We suggest a new mass generation mechanism for gauge fields. The quantum field theory constructed in this paper is nonabelian, gauge invariant and asymptotically free.

2000 Mathematics Subject Classification: 81T13.

Keywords and phrases: Yang-Mills field, mass generation.

It is well known that the naive mass generation mechanism can not be applied in case of the nonabelian Yang-Mills theory. We recall that if $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the strength tensor of the abelian $U(1)$ gauge field A_μ , $\mu = 0, 1, 2, 3$, and S is the usual Maxwell action,

$$S = - \int \frac{1}{4} F_{\mu\nu} F^{\mu\nu} d^4x, \quad (1)$$

then one can easily construct the action S_m^0 generating the corresponding massive theory (see, for instance, [8], Sect. 3.2.3),

$$S_m^0 = \int \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{m^2}{2} A_\mu A^\mu \right) d^4x. \quad (2)$$

Here and thereafter we use the standard convention about summations and lowering tensor indexes with the help of the standard metric $g_{\mu\nu}$ of the Minkowski space, $g_{00} = 1$, $g_{ii} = -1$ for $i = 1, 2, 3$, and $g_{ij} = 0$ for $i \neq j$.

In contrast to (1) action (2) is not invariant under gauge transformations

$$A_\mu \mapsto A_\mu + \partial_\mu \varphi, \quad (3)$$

where φ is any scalar function on the Minkowski space. However, if $m \neq 0$ the equations of motion generated by action (2),

$$\partial^\mu F_{\mu\nu} + m^2 A_\nu = 0, \quad (4)$$

imply the Lorentz gauge fixing condition

$$\partial^\mu A_\mu = 0. \quad (5)$$

Condition (5) is obtained by taking the divergence of equation (4).

Since action (2) is quadratic in A_μ the field theory associated to (2) can be easily quantized, the key point in the quantization procedure being the gauge fixing condition (5) which automatically appears in the massive case.

If G is a compact simple Lie group the construction of the massive quantum vector field theory outlined above can not be carried over in case of the Yang-Mills field associated to G . Formally the action similar to (2) can be written down. But the new action containing selfinteraction terms turns out to be non-renormalizable (see [8], Sect. 12.5.2). The main obstruction to renormalizability in the nonabelian case is nonlinearity of the corresponding gauge action. This leads, in turn, to the fact that the gauge fixing condition similar to (5) does not automatically appear in the nonabelian case.

Usually one overcomes this difficulty by using spontaneous symmetry breaking and the Higgs mechanism [5, 6]. But as a byproduct of that mechanism one gets extra unwanted boson fields.

In order to avoid the Higgs mechanism for constructing a massive nonabelian gauge quantum theory we start by slightly modifying action (2) in such a way that the new action S_m for the massive field is invariant under gauge transformations (3). We introduce the new action S_m by the following formula

$$S_m = \int \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{m^2}{2} (PA)_\mu (PA)^\mu \right) d^4x. \quad (6)$$

Here

$$(PA)_\mu = A_\mu - \partial_\mu \square^{-1} \partial^\nu A_\nu \quad (7)$$

is the orthogonal projection of the field A_μ onto the kernel of the divergence operator $\partial \cdot A = \partial^\mu A_\mu$, with respect to the scalar product $\langle \cdot, \cdot \rangle$ of vector fields,

$$\langle A, B \rangle = \int A_\mu(x) B^\mu(x) d^4x,$$

and $\square = \partial^\mu \partial_\mu$ is the d'Alembert operator. In formula (7) it is assumed that an inverse to the d'Alembert operator acting on the image of $\square$ is fixed. This

can be achieved, for instance, by choosing boundary conditions for large $x_0 = t$. The corresponding boundary conditions should be also imposed on gauge transformations φ in formula (3) to insure that each φ belongs to the image of $\square^{-1}$ and the identity

$$\square^{-1}\square\varphi = \varphi \quad (8)$$

holds. The particular choice of the operator $\square^{-1}$ is actually not important. In this paper we shall assume that all inverse d'Alembert operators are of Feynman type.

Observe that $(PA)_\mu$ is the gauge transform (3) of A_μ for $\varphi = -\square^{-1}\partial^\nu A_\nu$.

By definition the quantity $(PA)_\mu$ is invariant under gauge transformations (3). Indeed, if $A_\mu \mapsto A_\mu + \partial_\mu\varphi$ then

$$(PA)_\mu \mapsto A_\mu + \partial_\mu\varphi - \partial_\mu\square^{-1}\partial^\nu(A_\nu + \partial_\nu\varphi) = (PA)_\mu.$$

$(PA)_\mu$ is actually the gauge invariant part of A_μ . Moreover, $(PA)_\mu$ satisfies the Lorentz condition (5),

$$\partial^\mu(PA)_\mu = \partial^\mu A_\mu - \partial^\mu\partial_\mu\square^{-1}\partial^\nu A_\nu = 0. \quad (9)$$

Note that $(PA)_\mu$ linearly depends on A_μ since the gauge action is linear in the abelian case.

Since $(PA)_\mu$ is invariant under gauge transformations action (6) is also gauge invariant.

Observe that the lagrangian density associated to (6) is not local. From the first sight this is a serious obstruction for using action (6) in quantum field theory. However, if

$$\partial^\mu A_\mu = 0$$

one has by definition (7)

$$(PA)_\mu = A_\mu - \partial_\mu\square^{-1}\partial^\nu A_\nu = A_\mu.$$

Therefore after imposing gauge condition (5) action (6) coincides with action (2) and, hence, generates the same equations of motion as (2) modulo gauge transformations. We conclude that action (6) can be quantized similarly to (2), and the corresponding Green functions for *local gauge invariant* observables constructed with the help of (6) satisfy the locality condition.

From the above discussion we infer that actions (2) and (6) are physically equivalent and generate the same quantum field theory. In this sense action (6) is a trivial generalization of (2). But a remarkable property of the former one is that it has a *gauge invariant renormalizable* counterpart in the nonabelian case!

Formula (6) is not yet the most suitable for generalizing to the nonabelian case since it contains the operator P which is related to the explicit form of the gauge action (3). In order to exclude the operator P from formula (6) we

observe that since $(PA)_\mu$ is a gauge transform of A_μ , and the strength tensor is gauge invariant, one also has

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu = \partial_\mu (PA)_\nu - \partial_\nu (PA)_\mu. \quad (10)$$

Now using (9), (10) and integrating by parts we obtain the following formula

$$-\int \frac{1}{2} (\Box^{-1} F_{\mu\nu}) F^{\mu\nu} d^4x = \int (PA)_\mu (PA)^\mu d^4x.$$

The last relation allows to rewrite action (6) in an alternative form,

$$S_m = -\frac{1}{4} \int (F_{\mu\nu} F^{\mu\nu} + m^2 (\Box^{-1} F_{\mu\nu}) F^{\mu\nu}) d^4x. \quad (11)$$

The linearity of the gauge action in the abelian case is actually not crucial for formula (11) since it does not explicitly contain the operator P .

Now it is easy to write down a nonabelian counterpart of action (11). First we fix the notation as in [8]. Let G be a compact simple Lie group, $\mathfrak{g}$ its Lie algebra with the commutator denoted by $[\cdot, \cdot]$. We fix a nondegenerate invariant under the adjoint action scalar product on $\mathfrak{g}$ denoted by tr (for instance, one can take the trace of the composition of the elements of $\mathfrak{g}$ acting in the adjoint representation). Let t^a , $a = 1, \dots, \dim \mathfrak{g}$ be a linear basis of $\mathfrak{g}$ normalized in such a way that $\text{tr}(t^a t^b) = -\frac{1}{2} \delta^{ab}$.

We denote by A_μ the $\mathfrak{g}$ -valued gauge field (connection on the Minkowski space),

$$A_\mu = A_\mu^a t^a.$$

Let D_μ be the associated covariant derivative,

$$D_\mu = \partial_\mu - g A_\mu,$$

where g is a coupling constant, and $F_{\mu\nu}$ the strength tensor (curvature) of A_μ ,

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - g[A_\mu, A_\nu].$$

We shall also need a covariant d'Alambert operator $\Box_A$ associated to the gauge field A_μ ,

$$\Box_A = D_\mu D^\mu.$$

The covariant d'Alambert operator can be applied to any tensor field defined on the Minkowski space and taking values in a representation space of the Lie algebra $\mathfrak{g}$, the $\mathfrak{g}$ -valued gauge field A_μ acts on the tensor field according to that representation. Note that the operator $\Box_A$ is scalar, i.e. it does not change types of tensors.

Finally recall that the gauge group of G -valued functions $g(x)$ defined on the Minkowski space acts on the gauge field A_μ by

$$A_\mu \mapsto \frac{1}{g} (\partial_\mu g) g^{-1} + g A_\mu g^{-1}. \quad (12)$$

The corresponding transformation laws for the covariant derivative and the strength tensor are

$$D_\mu \mapsto g D_\mu g^{-1}, \quad (13)$$

$$F_{\mu\nu} \mapsto g F_{\mu\nu} g^{-1}. \quad (14)$$

Formula (13) implies that the covariant d'Alembert operator is transformed under gauge action (12) as follows

$$\square_A \mapsto g \square_A g^{-1}. \quad (15)$$

In the last formula we assume that the gauge group acts on tensor fields according to the representation of the group G induced by that of the Lie algebra $\mathfrak{g}$.

From formulas (14) and (15) it follows that the natural nonabelian analogue of action (11) invariant under gauge action (12) is

$$S_m^G = \frac{1}{2} \int \text{tr} (F_{\mu\nu} F^{\mu\nu} + m^2 (\square_A^{-1} F_{\mu\nu}) F^{\mu\nu}) d^4x, \quad (16)$$

where in the expression $(\square_A^{-1} F_{\mu\nu})$ we assume that $F_{\mu\nu}$ is in the adjoint representation of $\mathfrak{g}$ and the scalar operator $\square_A^{-1}$ acts on $F_{\mu\nu}$ componentwise. The first term in (16) is simply the action of the Yang-Mills field.

Now a new problem immediately appears. Due to nonlinearity of the gauge action in the nonabelian case it is impossible to make the lagrangian density associated to action (16) local by imposing a gauge fixing condition. However, the appearance of nonlocal quantities in quantum field theory is not a new fact. One should recall, first of all, the Faddeev-Popov determinant [3] that can be made local using additional anticommuting ghost fields. In this paper we are going to apply a similar trick to action (16).

Similarly to the case of pure Yang-Mills field (see [8], Sect. 12.2; [4], Ch. 3, §3) let us first formally write down the expression for the generating functional $Z(J)$ of the Green functions associated to action (16) via a Feynman path integral,

$$Z(J) = \int \mathcal{D}(A) \mathcal{D}(\eta) \mathcal{D}(\bar{\eta}) \exp \{ i \int [\text{tr} (\frac{1}{2} F_{\mu\nu} F^{\mu\nu} + \frac{m^2}{2} (\square_A^{-1} F_{\mu\nu}) F^{\mu\nu} + 2\bar{\eta}(\mathcal{M}\eta) - 2J^\mu A_\mu) + \mathcal{F}(A)] d^4x \}, \quad (17)$$

where J^μ is the source for A_μ taking values in the Lie algebra $\mathfrak{g}$, $\mathcal{F}(A)$ is a gauge fixing term, $\mathcal{M}$ the corresponding Faddeev-Popov operator, $\bar{\eta}, \eta$ are the anticommuting scalar ghost fields taking values in the adjoint representation of $\mathfrak{g}$ (Faddeev-Popov ghosts). Here and later we always assume that the measure $\mathcal{D}$ in Feynman path integrals is properly normalized.

The key observation is that the nonlocal part of action (16) is the quadratic form of the operator $\square_A^{-1}$. Therefore to make the expression in the exponent in

formula (17) explicitly local it is natural to use a formula for Gaussian integrals relating the exponent of the quadratic form of the operator and the exponent of the quadratic form of the inverse operator. For operator K acting on scalar fields that formula takes the following form ([4], Ch. 2, §6)

$$\begin{aligned} \exp\left\{\frac{i}{2} \int (K^{-1}\varphi)\varphi d^4x\right\} &= \\ &= \det(K)^{\frac{1}{2}} \int \mathcal{D}(\psi) \exp\left\{i \int \left(-\frac{1}{2}(K\psi)\psi + \varphi\psi\right) d^4x\right\}. \end{aligned} \quad (18)$$

Since in formula (17) the operator $\square_A^{-1}$ acts on the skew-symmetric (2,0)-type tensor field $F_{\mu\nu}$, in order to apply a formula similar to (18) to the expression

$$\exp\left\{i \int \text{tr}\left(\frac{m^2}{2}(\square_A^{-1}F_{\mu\nu})F^{\mu\nu}\right) d^4x\right\} \quad (19)$$

one has to introduce a new skew-symmetric (2,0)-type tensor field $\Phi_{\mu\nu}$ (bosonic ghost) with values in the adjoint representation of the Lie algebra $\mathfrak{g}$.

Using the ghost field $\Phi_{\mu\nu}$ and formula (18) we can rewrite expression (19) in an alternative form,

$$\begin{aligned} \exp\left\{i \int \text{tr}\left(\frac{m^2}{2}(\square_A^{-1}F_{\mu\nu})F^{\mu\nu}\right) d^4x\right\} &= \\ &= (\det \square_A|_{\Omega^3(\mathfrak{g})})^{\frac{1}{2}} \int \mathcal{D}(\Phi) \exp\left\{i \int \text{tr}\left(-\frac{1}{8}(\square_A\Phi_{\mu\nu})\Phi^{\mu\nu} + \frac{m}{2}\Phi_{\mu\nu}F^{\mu\nu}\right) d^4x\right\}, \end{aligned} \quad (20)$$

where the operator $\square_A$ is assumed to act componentwise in the space $\Omega^2(\mathfrak{g})$ of skew-symmetric (2,0)-type tensors with values in the adjoint representation of $\mathfrak{g}$. The numerical factors in the r.h.s. of the last formula appear since the Feynman path integral in (20) is only taken with respect to the linearly independent components $\Phi_{\mu\nu}$, $\mu < \nu$ of the skew-symmetric tensor $\Phi_{\mu\nu}$.

The r.h.s. of (20) still contains a nonlocal term, $(\det \square_A|_{\Omega^2(\mathfrak{g})})^{\frac{1}{2}}$. But since $\Omega^2(\mathfrak{g})$ is the direct sum of six copies of the space $\Omega^0(\mathfrak{g})$ of scalar fields with values in the adjoint representation of $\mathfrak{g}$ and the operator $\square_A$ acts componentwise in $\Omega^2(\mathfrak{g})$ the determinant $\det \square_A|_{\Omega^2(\mathfrak{g})}$ is equal to $(\det \square_A|_{\Omega^0(\mathfrak{g})})^6$, and

$$(\det \square_A|_{\Omega^2(\mathfrak{g})})^{\frac{1}{2}} = (\det \square_A|_{\Omega^0(\mathfrak{g})})^3.$$

Therefore recalling the Faddeev-Popov trick (see [4], Ch. 3, §3; [8], Sect. 12.2.2) and introducing six scalar anticommuting ghost fields $\eta_i, \bar{\eta}_i$, $i = 1, 2, 3$ taking values in the adjoint representation of $\mathfrak{g}$ we can express the square root of the determinant, $(\det \square_A|_{\Omega^2(\mathfrak{g})})^{\frac{1}{2}}$, in the following form

$$\begin{aligned} (\det \square_A|_{\Omega^2(\mathfrak{g})})^{\frac{1}{2}} &= (\det \square_A|_{\Omega^0(\mathfrak{g})})^3 = \\ &= \int \prod_{i=1}^3 \mathcal{D}(\eta_i) \mathcal{D}(\bar{\eta}_i) \exp\left\{i \int 2\text{tr}\left(\sum_{i=1}^3 \bar{\eta}_i (\square_A \eta_i)\right) d^4x\right\}. \end{aligned} \quad (21)$$

Now substituting (20) and (21) into (17) we derive an expression for the generating function $Z(J)$ that only contains local interaction terms and suitable for developing perturbation theory,

$$\begin{aligned}
Z(J) = & \int \mathcal{D}(A) \mathcal{D}(\Phi) \mathcal{D}(\eta) \mathcal{D}(\bar{\eta}) \prod_{i=1}^3 \mathcal{D}(\eta_i) \mathcal{D}(\bar{\eta}_i) \times \\
& \times \exp \left\{ i \int [\text{tr} \left(\frac{1}{2} F_{\mu\nu} F^{\mu\nu} - \frac{1}{8} (\Box_A \Phi_{\mu\nu}) \Phi^{\mu\nu} + \frac{m}{2} \Phi_{\mu\nu} F^{\mu\nu} + \right. \right. \\
& \left. \left. + 2\bar{\eta}(\mathcal{M}\eta) + 2 \sum_{i=1}^3 \bar{\eta}_i (\Box_A \eta_i) - 2J^\mu A_\mu \right) + \mathcal{F}(A)] d^4x \right\}. \quad (22)
\end{aligned}$$

Formula (22) looks a little bit complicated. But it is equivalent to the compact expression (17). Note that when deriving (22) starting from (17) we only used tricks with Gaussian Feynman path integrals. From (17) it also follows that when $m = 0$ the generating function $Z(J)$ is reduced to that of the quantized Yang-Mills theory.

Now we can discuss renormalization of the lagrangian in formula (22). In order to do that we have to first fix a gauge, i.e. we have to choose a gauge fixing term $\mathcal{F}(A)$ and specify the Faddeev-Popov operator $\mathcal{M}$.

We shall see in a moment that for our purposes the most convenient choice of $\mathcal{F}(A)$ is

$$\mathcal{F}(A) = \text{tr}(\partial^\mu A_\mu \partial^\nu A_\nu + m^2 \partial^\mu A_\mu (\Box^{-1} \partial^\nu A_\nu)). \quad (23)$$

Since the r.h.s. of (23) only depends on the divergence $\partial^\mu A_\mu$ of A_μ the corresponding Faddeev-Popov operator has the same form as in case of the Lorentz gauge (see [4], Ch. 3, §3),

$$\mathcal{M} = \partial^\mu D_\mu. \quad (24)$$

Note that gauge fixing condition (23) contains a nonlocal term. But since the nonlocal term only depends on the longitudinal component of the gauge field A_μ the nonlocal part in (23) does not generate extra nonlocal terms in the expressions for gauge invariant physical observables (see [4], Ch. 3, §3 for discussion of a similar situation in case of pure Yang-Mills field).

Finally combining (22), (23) and (24) we arrive at the following formula for $Z(J)$

$$\begin{aligned}
Z(J) = & \int \mathcal{D}(A) \mathcal{D}(\Phi) \mathcal{D}(\eta) \mathcal{D}(\bar{\eta}) \prod_{i=1}^3 \mathcal{D}(\eta_i) \mathcal{D}(\bar{\eta}_i) \times \\
& \times \exp \left\{ i \int \text{tr} \left(\frac{1}{2} F_{\mu\nu} F^{\mu\nu} - \frac{1}{8} (\Box_A \Phi_{\mu\nu}) \Phi^{\mu\nu} + \frac{m}{2} \Phi_{\mu\nu} F^{\mu\nu} - 2J^\mu A_\mu + \right. \right. \\
& \left. \left. + 2\bar{\eta}(\partial^\mu D_\mu \eta) + 2 \sum_{i=1}^3 \bar{\eta}_i (\Box_A \eta_i) + \partial^\mu A_\mu \partial^\nu A_\nu + m^2 \partial^\mu A_\mu (\Box^{-1} \partial^\nu A_\nu) \right) d^4x \right\}. \quad (25)
\end{aligned}$$

By simple dimensional counting the quantum field theory with generating function (25) is renormalizable. Moreover, using dimensional regularization and the BRST technique one can show, similarly to the case of the Yang-Mills field (see [8]), that gauge invariance is preserved by the renormalization. The details of those proofs and of other calculations that we are going to discuss can be found in [14].

The particular choice (23) of the gauge fixing term is convenient since in this case the free propagator $D_{\mu\nu}^{ab}$ of the gauge field A_μ has a very simple form,

$$D_{\mu\nu}^{ab} = i\delta^{ab} \frac{g_{\mu\nu}}{\square + m^2}.$$

Therefore the gauge field A_μ becomes massive. However, the theory may also contain some massless particles corresponding to the ghost fields.

The coupling constant renormalization calculated at the one-loop order using dimensional regularization coincides with that for the pure Yang-Mills field (see [8]),

$$g_0 = g\mu^{\frac{\epsilon}{2}} \left(1 - \frac{g^2}{16\pi^2} \frac{11}{6} C \frac{2}{\epsilon} \right), \quad (26)$$

where g_0 is the bare coupling constant, $\epsilon \rightarrow 0$ when the regularization is removed, μ is an arbitrary mass scale, and the constant C is defined by

$$C\delta_{ab} = \sum_{c,d} C_a^{cd} C_b^{cd}$$

with the help of the structure constants C_c^{ab} of the Lie algebra $\mathfrak{g}$, $[t^a, t^b] = C_c^{ab} t^c$.

It is easy to see that the coupling constant renormalization in the theory with generating function (25) calculated within the dimensional regularization framework coincides with the coupling constant renormalization for the pure Yang-Mills theory to all orders of perturbation theory. Indeed, by the results of [7] in the case of dimensional regularization the coupling constant renormalization is independent of the mass for dimensional reasons. Therefore the coupling constant renormalization in the theory with generating function (25) is the same as in the massless case, i.e. in case of the Yang-Mills field.

One can also calculate the mass renormalization at the one-loop order,

$$m_0 = m\mu^{\frac{\epsilon}{2}} \left(1 - \frac{g^2}{16\pi^2} \frac{11}{6} C \frac{2}{\epsilon} \right), \quad (27)$$

where m_0 is the bare mass.

The wave function renormalization at the one-loop order has the same form as in case of the Yang-Mills field,

$$Z_A = 1 + \frac{g^2}{16\pi^2} \frac{5}{3} C \frac{2}{\epsilon}. \quad (28)$$

From (26), (27) and (28) one can get the corresponding renormalization group coefficients at the one-loop order (see [13], Sect. 4.5),

$$\beta(g) = \mu \frac{\partial g}{\partial \mu} = -\frac{g^3}{16\pi^2} \frac{11}{3} C, \quad (29)$$

$$\gamma_m(g) = \mu \frac{\partial \ln m}{\partial \mu} = -\frac{g^2}{16\pi^2} \frac{11}{6} C, \quad (30)$$

$$\gamma_d(g) = \frac{1}{2} \mu \frac{\partial \ln Z_A}{\partial \mu} = -\frac{g^2}{16\pi^2} \frac{5}{3} C. \quad (31)$$

The Fourier transform $\Gamma(p_1, \dots, p_n, m, g, \mu)$ of any one-particle irreducible Green function with engineering dimension d is transformed under rescaling of the momenta as follows (see [13], Sect. 4.6)

$$\begin{aligned} \Gamma(sp_1, \dots, sp_n, m, g, \mu) &= \\ &= s^d \Gamma(p_1, \dots, p_n, m(s), g(s), \mu) \exp \left\{ -n \int_1^s \frac{ds'}{s'} \gamma_d(g(s')) \right\}, \end{aligned} \quad (32)$$

where $s > 0$ is a scale for the momenta. In equation (32) $g(s)$ and $m(s)$ are the solutions to the following equations

$$s \frac{\partial g(s)}{\partial s} = \beta(g(s)), \quad g(1) = g; \quad (33)$$

$$s \frac{\partial m(s)}{\partial s} = m(s)(\gamma_m(g(s)) - 1), \quad m(1) = m. \quad (34)$$

Applying standard renormalization group arguments (see, for instance, [8], [13]) we have $g(s) \rightarrow 0$ when $s \rightarrow \infty$ since the beta function $\beta(g)$ is negative in (33), and the theory with generating function (25) is asymptotically free.

Similarly, $m(s) \rightarrow 0$ when $s \rightarrow \infty$ since in that case $g(s) \rightarrow 0$ as we have just observed, and hence by (30) $\gamma_m(g(s)) - 1$ is negative in (34) for large s . Therefore the mass can be neglected for large momenta.

In conclusion we remark that the mass term in action (16) can be regarded as the result of a renormalization of the Yang-Mills action. Such possibility was discussed in [2]. However, the mass term can not be observed perturbatively at large momenta. We also note that there should be no serious problems with generalizing to our theory the program of rigorous construction of Green functions for the quantized Yang-Mills field outlined in [12].

One can also add to action (16) a fermionic part coupled to the gauge field A_μ in the minimal way.

Acknowledgements. When this work was completed the author has received information that the gauge invariant mass term that appears in formula (11) in the abelian case was considered in [10], and action (16) in the nonabelian

case was recently introduced and studied in [1]. Paper [1] also contains an extensive list of references on dynamical mass generation in the Yang-Mills theory.

Another action describing massive gauge fields was proposed in [11].

Prof. R. Jackiw also communicated to me that in case of three dimensions the mass generation mechanism similar to the one introduced in this paper was suggested in [9]. The author would like to thank him, David Dudal, Amitabha Lahiri and Jean-Luc Jacquot for pointing out the references quoted above.

References

- [1] Capri, M. A. L., Dudal, D., Gracey, J. A., Lemes, V. E. R., Sobreiro, R. F., Sorella, S. P., Verschelde, H., Study of the gauge invariant, nonlocal mass operator $\text{Tr} \int d^4x F_{\mu\nu} (D^2)^{-1} F_{\mu\nu}$ in Yang-Mills theories, *Phys. Rev. D* **72** (2005), 105016.
- [2] Faddeev, L. D., Mass in quantum Yang-Mills theory (comment on a Clay millenium problem), *Bull. Braz. Math. Soc., New Series* **33(2)** (2002), 201–212.
- [3] Faddeev, L. D., Popov V. N., Feynman diagrams for the Yang-Mills field, *Phys. Lett. B* **25** (1967), 29.
- [4] Faddeev, L. D., Slavnov, A. A., Gauge fields. Introduction to quantum theory, *Frontiers in Physics*, 83 Addison-Wesley (1991).
- [5] Higgs, P. W., Broken symmetries and the masses of gauge bosons, *Phys. Rev. Lett.* **13** (1964), 508–509.
- [6] Higgs, P. W., Spontaneous symmetry breakdown without massless bosons, spontaneous symmetry breakdown without massless bosons, *Phys. Rev. (2)* **145** (1966), 1156–1163.
- [7] 't Hooft, G., Dimensional regularization and the renormalization group, *Nucl. Phys. B* **61** (1973), 455–468.
- [8] Itzykson, C., Zuber, J.-B., Quantum Field Theory, McGraw-Hill (1980).
- [9] Jackiw, R., Pi, S.-Y., Seeking an even-parity mass term for 3-D gauge theory, *Phys. Lett. B* **403** (1997), 297.
- [10] Jacquot, J. L., Path integral regularization of QED by means of Stueckelberg fields, *Phys. Lett. B* **631** (2005) 83.
- [11] Lahiri, A., Generating Vector Boson Masses, preprint hep-th/9301060; The Dynamical Nonabelian Two-Form: BRST Quantization, *Phys. Rev. D* **55** (1997), 5045–5050; Renormalizability of the Dynamical Two-Form, *Phys. Rev. D* **63** (2001), 105002.

- [12] Magnen, J., Rivasseau, V., Sénéor, R., Construction of YM_4 with an infrared cutoff, *Comm. Math. Phys.* **155** (1993), 325–383.
- [13] Ramond, P., Field theory: a modern primer, Addison-Wesley (1989).
- [14] Sevostyanov, A., Nonlocal lagrangians and mass generation for gauge fields, preprint hep-th/0605051.